

Richard Katz, M.D.  
 Morton Schneider, M.D.  
 Steven Albert, M.D.  
 Alison Bender Haimes, M.D.  
 Stephen D. Greenberg, M.D.  
 Douglas R. DeCorato, M.D.  
 Gavin L. Duke, M.D.



Barbara H. Braffman, M.D.  
 Robert Ludwig, M.D.  
 Paul Choi, M.D.  
 Sean Herman, M.D.  
 Clyde Hershman, M.D.  
 Gwen Harris, M.D.

519 & 523 East 72nd Street • New York, NY 10021 • 430 East 59th Street, Sutton Place • New York, NY 10022  
 Tel: 212-288-1575 • Fax: 212-288-7616 • www.eastriverimaging.com

**XRAY**

<b>Esophagram and/or GI and/or Small Bowel</b>	Nothing to eat or drink 12 hours prior to exam
<b>Barium Enema and Air Contrast</b>	<p>Clear liquid diet for 24 hours prior to exam. Liquids include clear juices such as apple, cranberry &amp; grape, clear soups, jello, coffee or tea. No milk products or carbonated beverages.</p> <p><b>AT 8:00 P.M.</b> Night before - Drink one 10oz. bottle of citrate of magnesia.</p> <p><b>Beginning at 8 P.M. until bedtime</b>, drink four 8 oz. glasses of water.</p> <p><b>AT 10.P.M.</b> Night before - Take four Dulcolax pills with water. On the morning of the exam you may have any liquid listed above.</p> <p><b>AFTER A BARIUM STUDY</b> take a laxative (3 tbsp. Milk of Magnesia)</p>
<b>Gallbladder (OCG)</b>	Call our office for preparation
<b>Intravenous Pyelogram (IVP)</b>	<p><b>AT 6 P.M.</b> Night before - Drink one 10 oz bottle of citrate of magnesia. You may have a moderate supper.</p> <p><b>Beginning at 8 p.m.</b> until bedtime drink four 8 oz glasses of water.</p> <p><b>Nothing to eat or drink</b> after midnight until your exam is completed.</p>
<b>Plain X-rays</b>	There is no preparation for these studies.